

Leukaemia Section

Short Communication

t(9;10)(q34;q22) ZMIZ1/ABL1

Jean-Loup Huret

Genetics, Dept Medical Information, University of Poitiers, CHU Poitiers Hospital, F-86021 Poitiers, France (JLH)

Published in Atlas Database: December 2013

Online updated version : <http://AtlasGeneticsOncology.org/Anomalies/t0910q34q22ID1520.html>
DOI: 10.4267/2042/53974

This work is licensed under a Creative Commons Attribution-Noncommercial-No Derivative Works 2.0 France Licence.
© 2014 Atlas of Genetics and Cytogenetics in Oncology and Haematology

Abstract

Short communication on t(9;10)(q34;q22) ZMIZ1/ABL1, with data on clinics, and the genes implicated.

Clinics and pathology

Note

Only five cases to date have been described with a t(9;10)(q34;q22).

Two cases of acute myeloid leukemia (AML) are herein excluded from the study, because they are likely to represent another entity, with perhaps different genes involved.

These are two patients aged 44 years (1 female and 1 male), one with a M1-AML, and the other one with a M3-AML and the classical t(15;17)(q22;q21), both with a complex karyotype and no further data (Sato et al., 1995; Schoch et al., 1996).

Disease

B-cell acute lymphoblastic leukemia (B-ALL).

Phenotype/cell stem origin

One case was phenotyped as a CD10+ B-ALL.

Epidemiology

Three cases to date, 1 male / 2 female patients, all are paediatric cases so far, one was a 1.5 year old baby (Cooley et al., 2007; Rabin et al., 2008; Soler et al., 2008).

Prognosis

Scarce data, no follow-up; one case was noted as being in complete remission.

Cytogenetics

Cytogenetics morphological

Additional abnormalities were found in the three cases (complex karyotype in two cases). A del(13q) was found in one case.

Genes involved and proteins

ABL1

Location


9q34.12

Protein

1130-1143 amino acids; ABL1 is composed from N-term to C-term of a CAP, a SRC homology 3 (SH3), SH2 and SH1 tyrosine kinase domains, a conserved PXXP motifs to mediate protein-protein interactions, three nuclear localization signal (NLS) motifs, a DNA-binding domain, a globular (G).actin-binding domain, one nuclear export signal (NES), and a conserved filamentous actin-binding domain. Regulates endocytosis, epithelial-to-mesenchymal transition, cell polarity, adhesion, migration and invasion; in response to cellular stress, including DNA damage, ABL1 induces growth arrest and/or apoptosis (review in Greuber et al., 2013).

Somatic mutations

Apart from the well known role of ABL1 in leukemias, ABL1 has been found overexpressed in a small percentage of breast carcinomas, lung squamous cell carcinomas, and uterine corpus carcinomas.



1812 amino acids (aa): the aa 708 of ZMIZ1 is fused to aa 26 ABL1

ZMIZ1: Pro: proline-rich; AR: binding region for AR; N: nuclear localization signal; Miz domain;

TAD: transcriptional activation domain

ABL1: C: CAP; 3: SH3; 2: SH2; 1: SH1; P: PXXP; N: nuclear localization signals; DNA: DNA-binding;

Actin: actin-binding; n: nuclear export signal

t(9;10)(q34;q22) ZMIZ1/ABL1

Jean Loup Huret 2013

© AtlasGeneticsOncology

t(9;10)(q34;q22) ZMIZ1/ABL1 protein.

ZMIZ1

Location

10q22.3

Protein

1067 amino acids (aa). ZMIZ1 is composed from N-term to C-term of a proline-rich region (aa 363-555 (334-555 according to Swiss-Prot)), a nuclear localization signal sequence (aa 691-713), a Miz domain (aa 738-790 (Zn finger 727-804 according to Swiss-Prot)), and a proline-rich region (transcriptional activation domain) (aa 824-1067 ((867-1002 according to Swiss-Prot))).

The central region between aa 556 and 790 is the primary binding region for AR (Sharma et al., 2003). Sequence specific DNA binding protein. Transcriptional coactivator related to members of the proteoglycan inhibitor of activated STAT (PIAS) family.

Co-activator of AR (androgen receptor).

Increases ligand-dependent transcriptional activity of AR, SMAD3, and TP53. ZMIZ1 is able to enhance the sumoylation of AR.

ZMIZ1 cooperates with NOTCH1 to regulate the C-MYC pathway.

ZMIZ1 is overexpressed in a subset of T-ALL (Rakowski et al., 2013).

Result of the chromosomal anomaly

Hybrid gene

Note

A ZMIZ1/ABL1 hybrid gene was searched for and detected in one case to date (Soler et al., 2003).

Description

5' ZMIZ1-3'ABL1. The exon 14 of ZMIZ1 is fused to exon 2 of ABL1.

Fusion protein

Description

197 kDa, 1812 amino acids fusion protein, containing the N-term proline-rich domain of ZMIZ1 and most of ABL1.

Oncogenesis

ZMIZ1/ABL1 may encode a constitutively activated tyrosine kinase.

References

Sato Y, Suto Y, Pietenpol J, Golub TR, Gilliland DG, Davis EM, Le Beau MM, Roberts JM, Vogelstein B, Rowley JD. TEL and KIP1 define the smallest region of deletions on 12p13 in hematopoietic malignancies. *Blood*. 1995 Aug 15;86(4):1525-33

Schoch C, Haase D, Haeflacher T, Freund M, Link H, Lengfelder E, Löffler H, Büchner T, Fonatsch C. Incidence and implication of additional chromosome aberrations in acute promyelocytic leukaemia with translocation t(15;17)(q22;q21): a report on 50 patients. *Br J Haematol*. 1996 Sep;94(3):493-500

Sharma M, Li X, Wang Y, Zarnegar M, Huang CY, Palvimo JJ, Lim B, Sun Z. hZimp10 is an androgen receptor co-activator and forms a complex with SUMO-1 at replication foci. *EMBO J*. 2003 Nov 17;22(22):6101-14

Cooley LD, Chenevert S, Shuster JJ, Johnston DA, Mahoney DH, Carroll AJ, Devidas M, Linda SB, Lauer SJ, Camitta BM. Prognostic significance of cytogenetically detected chromosome 21 anomalies in childhood acute lymphoblastic leukemia: a Pediatric Oncology Group study. *Cancer Genet Cytogenet*. 2007 Jun;175(2):117-24

Rabin KR, Man TK, Yu A, Folsom MR, Zhao YJ, Rao PH, Plon SE, Naeem RC. Clinical utility of array comparative

genomic hybridization for detection of chromosomal abnormalities in pediatric acute lymphoblastic leukemia. *Pediatr Blood Cancer*. 2008 Aug;51(2):171-7

Soler G, Radford-Weiss I, Ben-Abdelali R, Mahlaoui N, Ponceau JF, Macintyre EA, Vekemans M, Bernard OA, Romana SP. Fusion of ZMIZ1 to ABL1 in a B-cell acute lymphoblastic leukaemia with a *t(9;10)(q34;q22.3)* translocation. *Leukemia*. 2008 Jun;22(6):1278-80

Greuber EK, Smith-Pearson P, Wang J, Pendergast AM. Role of ABL family kinases in cancer: from leukaemia to solid tumours. *Nat Rev Cancer*. 2013 Aug;13(8):559-71

Rakowski LA, Garagiola DD, Li CM, Decker M, Caruso S,

Jones M, Kuick R, Cierpicki T, Maillard I, Chiang MY. Convergence of the ZMIZ1 and NOTCH1 pathways at C-MYC in acute T lymphoblastic leukemias. *Cancer Res*. 2013 Jan 15;73(2):930-41

Rogers LM, Riordan JD, Swick BL, Meyerholz DK, Dupuy AJ. Ectopic expression of Zmiz1 induces cutaneous squamous cell malignancies in a mouse model of cancer. *J Invest Dermatol*. 2013 Jul;133(7):1863-9

This article should be referenced as such:

Huret JL. *t(9;10)(q34;q22) ZMIZ1/ABL1*. *Atlas Genet Cytogenet Oncol Haematol*. 2014; 18(7):518-520.
